

Cultural Responsiveness in Trauma-Informed Care for Immigrant and First-Generation Populations in Urban Behavioral Health Settings

Sylvia Stella Favour Peteranaba^{1*}, Helen Ekwi Osinem², Florence Eribenne³, Olakunle Saheed Soyege⁴

¹Independent Researcher, Favourpeteranaba@gmail.com

²Jos University Teaching Hospital (JUTH),

Nursing Services Department, Jos, Nigeria helenosinem@gmail.com

³Bay Cove Human Services, Boston, MA, USA, feribenne1980@gmail.com

⁴Independent Researcher, Pretoria, south Africa, ossoyege@gmail.com

*Corresponding Author: Sylvia Stella Favour

Peteranaba, Favourpeteranaba@gmail.com

DOI: [10.56201/ijhpr.v8.no4.2023.pg154.174](https://doi.org/10.56201/ijhpr.v8.no4.2023.pg154.174)

Abstract

This review examines how culturally responsive, trauma-informed behavioral health can be strengthened for immigrant and first-generation populations in urban settings. It synthesizes multidisciplinary literature published through 2023 on migration-related adversity, cultural interpretations of trauma, intergenerational stress, language access, family and community influences, institutional trust, workforce competence, and organizational readiness. The review adopts a critical narrative approach, integrating evidence from global, African, and Nigerian contexts to assess how cultural and structural conditions shape engagement, assessment, treatment, and recovery. The findings indicate that trauma among immigrant and first-generation populations cannot be understood adequately through standardized clinical frameworks alone. Cultural identity, migration history, socioeconomic position, family expectations, racialization, language, and legal or institutional experiences influence how distress is expressed, interpreted, and addressed. Culturally responsive assessment requires cultural humility, context-sensitive measurement, recognition of idioms of distress, and careful attention to post-migration stressors. Language-concordant care and qualified interpretation are essential to therapeutic communication, while family, community, spiritual, and indigenous healing resources may strengthen recovery when incorporated safely and collaboratively. The review further identifies institutional mistrust, discrimination, financial barriers, fragmented services, and insufficient workforce preparation as persistent obstacles to equitable care. The study concludes that trauma-informed practice becomes genuinely responsive only when cultural meaning and structural inequity are embedded within clinical and organizational decision-making. It recommends sustained investment in interpreter services, culturally informed workforce development, reflective supervision, community partnerships, disparity monitoring, and policies that reduce coercive and administrative barriers. Future research should prioritize longitudinal, community-engaged, and culturally validated studies across diverse immigrant

generations and urban populations. Particular attention should be given to examining intersectional and intergenerational pathways to resilience.

Keywords: *cultural responsiveness; trauma-informed care; immigrant populations; first-generation populations; urban behavioral health; cultural humility.*

1. Introduction

1.1 Migration, Trauma, and Urban Behavioral Health

Migration is not intrinsically pathological, yet the conditions surrounding movement, settlement, and incorporation can expose immigrants and first-generation populations to layered forms of psychological strain. Pre-migration adversity may include political violence, persecution, community disruption, bereavement, or prolonged insecurity, while migration journeys can involve uncertainty, family separation, and loss of familiar social structures. After arrival, these experiences often intersect with housing instability, precarious employment, discrimination, legal uncertainty, language barriers, and weakened social networks. Accordingly, trauma-informed behavioral health must understand migration as a longitudinal social process rather than a single event. Evidence from refugee populations demonstrates that psychiatric morbidity reflects both prior exposure to adversity and the conditions encountered after resettlement, with depression, anxiety, and post-traumatic stress representing prominent concerns across displaced populations (Blackmore et al., 2020). Post-migration environments are especially consequential in urban settings, where proximity to services does not necessarily produce meaningful access. Cities may offer specialized behavioral health systems, ethnic communities, employment opportunities, and culturally diverse institutions, but these advantages coexist with overcrowding, poverty, administrative complexity, xenophobia, and fragmented care. Hynie (2018) shows that social determinants such as income, housing, employment, social support, and discrimination can materially shape refugee mental health after migration. This perspective is critical because it shifts clinical attention beyond trauma histories toward the everyday environments in which recovery is expected to occur. The psychological burden of migration is also mediated by the accumulation of post-migration stressors. Unresolved asylum processes, restricted employment, family separation, acculturative demands, and diminished social status may sustain distress long after physical safety has been achieved (Li, Liddell & Nickerson, 2016). In Johannesburg, research with forced migrants similarly demonstrates how precarious urban living, unmet basic needs, and limited access to mental health support can intensify vulnerability even within a major metropolitan setting (Walker & Vearey, 2023). Such findings are particularly relevant to urban behavioral health systems because they reveal why symptom-focused treatment alone may be insufficient when structural stressors remain active.

Cultural and generational position further complicate this landscape. Among African immigrants in the United States, gender, migration generation, socioeconomic conditions, and social experiences have been associated with differences in depression and anxiety, underscoring the heterogeneity concealed by broad labels such as “immigrant” or “African” (Escamilla & Saasa, 2020). Research among immigrant Latinos likewise indicates that exposure to political violence and psychosocial trauma intersects with service utilization, emphasizing that help-seeking is shaped by both trauma exposure and the social context of care (Fortuna, Porche & Alegría, 2008).

Nigeria provides an important contextual illustration of the need to attend to culturally situated psychological vulnerability. Sabboh (2023) identifies perceived stress, hopelessness, and self-esteem as interrelated psychological constructs among Nigerian young adults, reinforcing the importance of interpreting distress within locally meaningful social and cultural conditions. Evidence from university students in Lagos similarly links psychological factors with severe emotional vulnerability, highlighting the relevance of accessible and context-sensitive mental health support in densely populated urban environments (Yusuf & Sabboh, 2021). For immigrant and first-generation populations, these observations support an approach to urban behavioral health that recognizes pre-migration experiences, post-migration conditions, cultural meanings, generational adaptation, and structural inequalities as interacting determinants of psychological wellbeing. Trauma-informed care in such settings therefore requires not merely awareness of prior adversity, but a sophisticated understanding of how migration reorganizes identity, belonging, social resources, exposure to stress, and pathways into care. Within metropolitan systems, this ecological understanding has implications: screening, referral, psychotherapy, crisis response, and community outreach should be coordinated so that language needs, migration histories, family structures, socioeconomic pressures, and culturally patterned expressions of distress are treated as meaningful. These factors should inform engagement, assessment, continuity, and interpretation across behavioral health services.

1.2 Cultural Contexts of Trauma and Recovery

Cultural context is integral to trauma-informed care because it shapes how adversity is interpreted, narrated, embodied, disclosed, and addressed. For immigrant and first-generation populations in urban behavioral health settings, trauma may be understood through family obligations, spiritual beliefs, collective histories, gender roles, community expectations, or culturally specific idioms of distress. Consequently, clinicians who interpret suffering exclusively through Western diagnostic categories risk overlooking meanings that determine whether clients recognise symptoms, accept treatment, or regard professional care as legitimate. Kirmayer et al. (2011) demonstrate that immigrant and refugee mental health assessment requires attention to cultural explanations of illness, migration experiences, family context, and differing patterns of help-seeking. Recovery is culturally mediated. Rather than representing an individual return to psychological equilibrium, healing may involve restoration of family relationships, social belonging, dignity, religious connection, or communal functioning. A culture-informed systems approach therefore complements trauma-informed principles by situating intervention within layered psychosocial supports and community resources, rather than limiting recovery to symptom reduction (Im, Rodriguez & Grumbine, 2021). This orientation is important in diverse cities, where multiple cultural traditions coexist within behavioral health systems that may privilege standardized models of care. Evidence from African practice illustrates why therapeutic protocols require contextual adaptation. Research involving EMDR therapists across five African countries identified culturally responsive modifications involving language, collective forms of self-understanding, expressions of emotion, religious practices, music, dance, and locally intelligible metaphors (Mbazzi et al., 2021). Such findings are relevant for African-origin immigrant communities because meanings may persist, evolve, or become more salient under conditions of displacement and acculturation.

Within Nigeria, family and economic structures demonstrate how psychosocial distress is embedded within cultural expectations. Changing breadwinner roles in Ibadan were associated

with marital, parenting, and economic pressures, illustrating how gendered responsibilities and socioeconomic disruption can alter family functioning and psychological wellbeing (Ezeani & Sabboh, 2021). Among adolescents in Delta State, socioeconomic status, peer influence, and self-esteem were examined as psycho-sociological determinants of behavioural outcomes (Balogun & Sabboh, 2021). For immigrant and first-generation clients, culturally responsive trauma care should explore not only what happened, but what the experience means within the client's family, community, identity, and moral world.

1.3 First-Generation Identity and Intergenerational Stress

First-generation identity is formed within a field in which young people negotiate family heritage, host-society expectations, and developmental demands for autonomy. In immigrant families, these pressures can become pronounced when children acquire language, social norms, and institutional knowledge more rapidly than parents. Such differences may redistribute authority within the household and produce acculturation gaps experienced not simply as disagreement, but as threats to continuity, belonging, and parental legitimacy. Research with migrant families in Australia shows that young people may become "contested sites of culture" as they reconcile integration with preservation of family values, thereby intensifying intergenerational strain (Renzaho et al., 2017). For families with histories of displacement, these tensions may coexist with unresolved trauma, loss, and altered social roles. Among resettled South Sudanese families in Australia, changes in cultural expectations, religion, gender relations, and perceptions of freedom were associated with intergenerational conflict and family stress, illustrating how migration can transform established systems of authority and identity (Pittaway et al., 2023). In urban behavioral health settings, clinicians must distinguish normative generational negotiation from distress compounded by displacement, discrimination, or prior adversity. Family processes remain central to psychological development beyond migration-specific research. Nigerian evidence indicates that parenting styles and effortful control are associated with adolescent mental health, underscoring the importance of relational environments in emotional regulation and adjustment (Oyarekua & Sabboh, 2021). Related findings from Kaduna identify parental socioeconomic conditions and parenting patterns as psychosocial correlates of child abuse, demonstrating how family stress can be intensified by structural and interpersonal pressures (Sabbah & Yusuff, 2021). Parental influence also extends into identity-forming decisions: involvement by parents was significantly associated with career decision-making among secondary-school students in Oyo State (Ezeani et al., 2023).

1.4 Review Scope, Objectives, and Questions

This review examines how cultural responsiveness can be meaningfully integrated into trauma-informed behavioral health care for immigrant and first-generation populations in urban settings. Its scope encompasses the interaction of migration-related adversity, cultural identity, intergenerational dynamics, language, family systems, community influences, and structural conditions that shape both psychological distress and pathways to recovery. Particular attention is given to the way's trauma may be interpreted, expressed, and treated differently across cultural contexts, and to the implications of these differences for assessment, engagement, service delivery, and continuity of care. The principal objective is to evaluate how trauma-informed principles can be strengthened through culturally responsive practice without reducing culture to a fixed set of traits or assumptions. The review therefore considers cultural humility,

individualized assessment, linguistic accessibility, family and community participation, and sensitivity to migration histories as central elements of effective care. It also examines how institutional barriers, mistrust, workforce limitations, and inequities in access may compromise the responsiveness of urban behavioral health systems. The analysis further emphasizes avoiding cultural stereotyping while recognizing resilience, social networks, spirituality, and collective resources within diverse communities.

Three guiding questions structure the review. First, how do cultural background, migration experience, and first-generation identity influence the understanding and expression of trauma? Second, what clinical and organizational adaptations improve the relevance, accessibility, and acceptability of trauma-informed services for culturally diverse populations? Third, what structural, workforce, and policy conditions are necessary to sustain equitable and culturally responsive care in urban behavioral health environments? By addressing these questions, the review seeks to clarify current knowledge, identify persistent gaps, and establish a coherent basis for practice improvement, service-system reform, and future research focused on culturally diverse urban populations.

2. Foundations of Culturally Responsive Trauma Care

2.1 Principles of Trauma-Informed Behavioral Health

Trauma-informed behavioral health is best understood as an organizational orientation that assumes trauma may shape how individuals perceive safety, authority, relationships, symptoms, and treatment environments. Its foundational task is therefore not to elicit detailed trauma histories indiscriminately, but to organize care so that services minimize avoidable harm while strengthening dignity, agency, trust, and recovery. For immigrant and first-generation populations, this orientation is especially important because experiences of displacement, political instability, discrimination, family separation, precarious legal status, or acculturative strain may coexist with cultural meanings of suffering that differ from psychiatric frameworks. Core trauma-informed principles emphasize safety, trustworthiness, collaboration, choice, empowerment, and recognition of the social contexts in which distress develops, requiring systems to move from asking what is wrong with a person toward understanding what experiences may have shaped current needs (Sweeney et al., 2016).

Safety is both physical and psychological. In urban behavioral health settings, routine procedures—locked environments, abrupt questioning, unfamiliar authority structures, coercive practices, or poor communication—may be experienced as threatening by people whose histories include persecution, interpersonal violence, institutional abuse, or forced migration. Trauma-informed practice consequently requires predictable processes, transparent explanations, respectful boundaries, and careful attention to environmental triggers. It also demands that professionals recognize the potential for services themselves to reproduce powerlessness. Evidence from mental-health nursing literature identifies sensitivity to trauma histories, autonomy, relationship-based practice, and avoidance of insensitive responses as important expressions of trauma-informed care, underscoring the ethical imperative to prevent retraumatization (Cleary & Hungerford, 2015).

Trust and collaboration are equally central because trauma can disrupt confidence in institutions and interpersonal relationships. Clinicians should explain assessment procedures, confidentiality, treatment options, and limits of professional authority in understandable terms, while allowing clients participation in decisions whenever clinically feasible. This is particularly significant for

immigrant populations who may encounter language discordance or uncertainty about whether behavioral health systems are connected to immigration, law enforcement, or other state institutions. Trauma-informed care also requires thoughtful screening: inquiry about adversity should occur only when there is a clinical purpose, adequate privacy, and a pathway for response. Literature synthesizing trauma-informed healthcare stresses sensitive disclosure practices, therapeutic provider-patient relationships, autonomy, multidisciplinary collaboration, and referral pathways as core mechanisms for reducing distress and improving engagement (Reeves, 2015). Cultural responsiveness strengthens these principles by recognizing that safety, trust, autonomy, and recovery are culturally interpreted rather than universally experienced in identical ways. A client's preference for family involvement, spiritual coping, indirect disclosure, same-gender clinicians, or collective decision-making may reflect cultural frameworks rather than resistance. Conversely, clinicians should not assume that cultural background determines individual preferences. Trauma-informed behavioral health therefore requires cultural humility: providers must examine their own assumptions, invite the client's explanatory model, and negotiate care without stereotyping. The South African experience demonstrates the importance of locating trauma within histories of violence, inequality, social adversity, and mental-health burden, illustrating why context-sensitive responses must connect individual symptoms to community and structural realities (Wyatt et al., 2017).

Empowerment and strengths-based care further require attention to resilience rather than defining people exclusively by traumatic exposure. Immigrant and first-generation clients may possess protective resources in family networks, multilingualism, faith communities, cultural identity, educational aspirations, and transnational relationships. Clinical assessment should identify these capacities alongside risk, while recognizing that psychosocial environments influence behavior and self-regulation. Nigerian evidence linking adolescent moral disengagement with psychosocial factors reinforces the importance of examining social and psychological contexts when interpreting behavior, rather than relying solely on individualized deficit explanations (Sabbah, 2021). In practice, trauma-informed behavioral health should integrate these principles across reception, assessment, crisis response, psychotherapy, medication management, referral, discharge planning, and community partnerships. Staff training, reflective supervision, interpreter access, culturally responsive policies, and mechanisms for service-user feedback are essential organizational conditions. When consistently operationalized, these principles create urban behavioral health environments capable of responding to trauma without reducing culturally diverse clients to diagnoses, migration histories, or presumed vulnerabilities, while preserving choice, relational safety, continuity, and culturally meaningful pathways toward recovery.

2.2 Culture, Identity, and Trauma Interpretation

Culture mediates the interpretation of trauma by shaping what counts as suffering, how distress is communicated, where causation is located, and which forms of help are considered legitimate. For immigrant and first-generation populations, these frameworks may coexist with biomedical language acquired through schools, workplaces, media, and health systems. Trauma therefore cannot be understood solely through standardized symptom categories; culturally situated meanings influence whether intrusive memories, somatic sensations, spiritual concerns, withdrawal, anger, or disrupted relationships are recognized as illness, adversity, moral injury, or hardship. Cultural concepts of distress provide a framework for examining these variations

without assuming psychiatric classifications exhaust meanings attached to suffering (Lewis-Fernández & Kirmayer, 2019).

Clinical interpretation becomes complex when migration relocates people across cultural systems. Among Somali refugees displaced in Kenya, culturally recognizable idioms offered meaningful ways of describing distress, causation, and social disruption, demonstrating that trauma narratives are embedded in collective histories and explanatory traditions (Im, Ferguson & Hunter, 2017). In urban behavioral health settings, such meanings may influence disclosure, stigma, family involvement, and willingness to accept interventions.

Identity further structures trauma interpretation because individuals occupy multiple social positions simultaneously. First-generation clients may negotiate parental expectations, peer norms, socioeconomic constraints, racialization, religious commitments, and host-society pressures while constructing a coherent sense of self. Nigerian research among university students indicates that peer approval seeking, socioeconomic status, and locus of control are associated with self-handicapping behaviour, illustrating how social evaluation and perceived agency can shape responses to demanding environments (Sabbah & Ojo, 2023). Related evidence from Ibadan identifies psychological and social correlates of academic buoyancy, highlighting the importance of resilience when individuals confront recurring pressures (Yusuf, Sabbah & Asiyanbi, 2023). For trauma-informed practice, cultural responsiveness therefore requires more than demographic matching or knowledge of customary beliefs. Clinicians need to elicit the client's explanation of what happened, what the experience signifies, why distress persists, and what recovery would look like within family and community life. Cross-cultural trauma assessment should incorporate idioms of distress, explanatory models, ongoing adversity, and cultural humility alongside conventional diagnostic evaluation (Patel & Hall, 2021). This approach reduces the risk of pathologizing culturally normative responses while avoiding the opposite error of attributing significant suffering to culture alone. In urban systems, identity should be treated as dynamic, relational, and shaped by migration, generation, context, and lived experience.

2.3 Intersectionality in Immigrant Behavioral Health

Intersectionality provides a framework for understanding immigrant behavioral health because migration status never operates independently of race, gender, socioeconomic position, age, language, religion, or generation. These social locations interact within institutions and neighborhoods, shaping exposure to adversity, access to resources, and meanings attached to psychological distress. Analyses that attribute immigrant mental health primarily to culture or acculturation can obscure structural processes such as racialization, exclusion, and restrictive policy environments that distribute health risks unequally (Viruell-Fuentes, Miranda & Abdulrahim, 2012). This perspective is particularly relevant in urban behavioral health systems, where diverse populations encounter the same services from unequal social positions. Population-based evidence from California demonstrates that psychological distress varies across combinations of immigrant generation, race, poverty, and gender, indicating that no single identity category adequately captures vulnerability or protection (Mancenido et al., 2020). For first-generation clients, these intersections may influence whether trauma is disclosed, how symptoms are interpreted, and whether professional care feels accessible, safe, or culturally legitimate.

Intersectionality also directs attention to post-migration discrimination as a behavioral-health determinant. Among African refugees and migrants in Durban, South Africa, discriminatory experiences, family separation, income, marital status, and timing of arrival were associated with anxiety, depression, and post-traumatic stress symptoms, revealing how migration-related distress is produced through interacting social and material conditions rather than migration alone (Thela et al., 2017). Such evidence is pertinent to trauma-informed care because social marginalization can remain an active source of threat after resettlement.

Generational position further complicates these patterns. Cross-national research among adolescents found that immigration background interacted with socioeconomic status and gender in shaping mental well-being, while national policy and social context influenced the magnitude of inequalities (Kern et al., 2020). These findings caution against treating “immigrant youth” as a homogeneous clinical category.

Within Nigeria, psychosocial research likewise demonstrates the simultaneous influence of institutional and interpersonal environments. School climate, self-efficacy, social media, and peer pressure were associated with aggressive behaviour among adolescents in Ibadan, illustrating how behavioral outcomes emerge from interacting domains of experience (Sabbah & Alausa, 2021). Applied to immigrant behavioral health, an intersectional trauma-informed approach should assess not only trauma exposure but also racialization, gendered expectations, socioeconomic insecurity, generational position, family relationships, and institutional barriers. Such assessment strengthens cultural responsiveness by locating distress within the overlapping systems that shape both vulnerability and resilience.

3. Clinical Adaptation for Diverse Urban Populations

3.1 Culturally Responsive Assessment and Engagement

Culturally responsive assessment in trauma-informed behavioral health requires more than selecting a standardized instrument and translating its items. For immigrant and first-generation populations, assessment is an interpretive and relational process in which clinicians must establish safety while eliciting the client’s account of distress, migration, identity, family relationships, and social conditions. The objective is not to replace diagnostic rigor with cultural description, but to improve validity by examining how symptoms are expressed, understood, and contextualized. The Cultural Formulation Interview offers a structured method for exploring cultural identity, explanatory models, stressors, supports, and preferences for care, helping clinicians integrate cultural information into psychiatric evaluation rather than treating culture as an optional supplement (Aggarwal et al., 2020).

Trauma assessment requires caution because direct questioning can reproduce feelings of exposure or powerlessness before trust develops. Immigrant clients may have experienced war, persecution, detention, interpersonal violence, discrimination, or dangerous migration journeys, while first-generation clients may be managing inherited family narratives and acculturative pressures. In Canadian mental-health services, providers working with immigrants and refugees identified inconsistencies in transcultural assessment and emphasized flexible, reflexive, trauma-informed approaches that allow cultural interpretations of mental illness to emerge within the clinical encounter (Wylie et al., 2018). Accordingly, clinicians should clarify confidentiality, explain why sensitive questions are asked, obtain consent where appropriate, and avoid pressing for detailed disclosure when information is unnecessary for immediate care.

Measurement validity is equally important. Screening tools developed in one population may not perform identically when language, symptom meaning, response styles, or cultural norms differ. Research comparing refugees with German residents found that depressive and anxiety symptoms did not function uniformly across groups, demonstrating the limitations of assuming measurement equivalence because an instrument has been translated (Schlechter et al., 2021). Evidence from a Zulu-speaking South African community similarly showed that adding a culture-specific trauma questionnaire to a conventional diagnostic tool elicited additional traumatic events and culturally specific attributions that the standard assessment alone did not capture (Madigoe et al., 2017). These findings support psychometric discipline alongside locally meaningful inquiry.

Nigerian evidence further illustrates why instruments should be evaluated within intended populations. Sabboh's development and validation of a revised self-esteem measure for Nigerian adolescents and youths shows that constructs require attention to wording, contextual relevance, reliability, and construct validity when applied across cultural settings (Sabbah, 2022). In immigrant behavioral health, similar care is needed when assessing trauma, depression, anxiety, resilience, or self-concept among clients whose cultural frames differ from populations on which instruments were standardized. Engagement begins during assessment rather than after it. Clients are more likely to participate meaningfully when clinicians demonstrate respect, acknowledge uncertainty, and invite correction rather than claiming authority over the client's experience. Cultural humility has been associated with stronger therapeutic alliance and improved psychotherapy processes, reinforcing the value of another-oriented stance characterized by openness and reduced assumptions of superiority (Hook et al., 2013). This is especially relevant for first-generation clients who may resist reduction to parental or host-society categories.

Assessment should distinguish historical trauma from ongoing threats. Survivors of torture and persecution may present with culturally mediated symptoms, somatic complaints, silence, mistrust, or fragmented narratives; culturally informed evaluation therefore requires attention to construct equivalence and the relationship between subjective meanings and observable clinical phenomena (Raghavan, 2019). Clinicians should document current stressors such as housing insecurity, family separation, discrimination, legal uncertainty, and social isolation because these conditions may sustain distress and limit participation.

At the service level, meaningful engagement depends on whether institutions adapt to communities rather than attributing nonattendance solely to client reluctance. Work with Somali refugee communities in the United States demonstrates the importance of partnership, co-learning, community ownership, and cultural humility in improving mental-health access (Abdi et al., 2022). Urban behavioral health systems should connect culturally responsive assessment with engagement strategies, including transparent communication, collaborative goal setting, appropriate interpreter use, family involvement when desired, continuity with trusted providers, and partnerships with community organizations. Such practices allow assessment to function not merely as diagnostic classification, but as the first stage of a respectful therapeutic relationship in which cultural meaning, trauma history, strengths, and present circumstances are integrated into individualized care.

3.2 Language, Interpretation, and Therapeutic Communication

Language is not merely a vehicle for transmitting clinical information; in trauma-informed behavioral health, it is central to how distress is narrated, interpreted, and therapeutically

contained. For immigrant and first-generation populations, linguistic discordance can affect disclosure, diagnostic precision, informed consent, alliance, and continuity. Clients may possess conversational proficiency in the dominant language while lacking vocabulary to describe intrusive memories, grief, bodily sensations, or culturally embedded meanings of suffering. A systematic review of psychiatric care found that assessment conducted in a non-primary language can distort mental-status evaluation, while professional interpreters can improve communication and reduce clinically consequential errors compared with ad hoc arrangements (Bauer & Alegría, 2010).

Interpreter-mediated encounters therefore require deliberate clinical structure. Professional interpreters should be briefed about therapeutic purpose, confidentiality, preferred interpretation mode, and sensitive material, while clinicians should address clients directly. Refugee mental-health research shows that interpreter's function within complex relational spaces involving language, culture, trauma narratives, trust, and role boundaries; their effectiveness depends on preparation, collaboration, and recognition of emotional demands associated with interpreting traumatic material (Fennig & Denov, 2021). These considerations are important when clients fear that disclosure may circulate within close-knit communities.

Evidence from South African psychiatric services illustrate risks created when systems rely on untrained bilingual staff. Psychiatrists reported miscommunication, role confusion, and concerns about adequacy of care when ad hoc interpreters were used, highlighting the need for formal language services in multilingual mental-health environments (Hagan et al., 2020). The issue extends beyond psychiatry: Nigerian evidence identifies language barriers as constraints on access, patient satisfaction, comprehension, and quality within a highly multilingual health system, reinforcing the importance of linguistic competence and responsive communication structures (Diri, 2023). Therapeutic communication must remain culturally responsive rather than mechanically translated. Literal equivalence does not guarantee conceptual equivalence; terms such as trauma, depression, recovery, family, spirituality, or safety may carry different meanings across linguistic communities. Clinicians should check understanding, invite clients to describe preferred terminology, and clarify culturally specific metaphors or idioms without pathologizing them. In urban behavioral health settings, language access should be embedded across intake, assessment, crisis intervention, psychotherapy, medication discussions, discharge planning, and follow-up. Consistent use of qualified interpreters, translated materials, language-concordant staff where feasible, and communication protocols can strengthen trust while reducing the risk that linguistic difference becomes a structural barrier to trauma-informed care.

3.3 Family, Community, and Indigenous Healing Practices

Family, community, and indigenous healing practices are central to culturally responsive trauma-informed care because recovery is often relational rather than exclusively individual. Among immigrant and first-generation populations, experiences of adversity may be interpreted through kinship obligations, spiritual beliefs, collective memory, and culturally sanctioned support. Clinical models that privilege private, individual psychotherapy can overlook resources that clients regard as essential to safety, identity, and restoration. A systematic review of family interventions for trauma-affected refugees found that family-based approaches can address relational disruption, communication, parenting, and resilience, while highlighting the need for stronger cultural adaptation and rigorous evaluation (Mak & Wieling, 2022).

Family involvement, however, should never be assumed to be universally beneficial or culturally required. Trauma-informed practice demands that clinicians establish whether the client wishes relatives to participate, assess potential conflict or coercion, and protect confidentiality and autonomy. Nigerian evidence linking parental bonding with psychological and behavioral outcomes among undergraduates illustrates the importance of close family relationships in shaping young people's adjustment and functioning (Sabboh & Olugbade, 2021). For first-generation clients, family engagement may be especially valuable when migration has altered authority, caregiving expectations, language roles, or intergenerational communication.

Community resources can similarly extend recovery beyond formal behavioral health services. Faith communities, cultural associations, elders, mutual-aid networks, and traditional healers may provide belonging, moral interpretation, practical support, and culturally recognizable pathways for responding to distress. In Ghana, Muslim traditional healers described explanatory models and treatment practices grounded in spiritual understandings of mental disorder, demonstrating why clinicians must understand local healing systems rather than dismiss them as peripheral to care (Kpobi & Swartz, 2019). Respectful engagement can improve dialogue while requiring attention to consent, safety, evidence, and human rights.

Indigenous healing also challenges behavioral health systems to recognize the consequences of colonization, historical trauma, and cultural disruption. Australian scholarship argues that cultural safety becomes more meaningful when trauma-informed practice acknowledges collective histories and supports community-defined processes of healing (Tujague & Ryan, 2021). For urban immigrant populations, the implication is not to transplant indigenous traditions indiscriminately, but to ask what healing practices, relationships, rituals, spiritual resources, or community institutions matter to each client. Culturally responsive services should create space for family participation, collaborate with trusted community resources where appropriate, and integrate traditional or spiritual practices when desired, lawful, and compatible with clinical safety.

4. Structural Barriers and Service-System Challenges

4.1 Access, Trust, and Institutional Mistrust

Access to trauma-informed behavioral health services is determined not only by whether services exist, but by whether immigrant and first-generation populations perceive them as reachable, legitimate, safe, and responsive. In urban environments, geographic proximity to clinics can coexist with barriers including cost, insurance restrictions, language discordance, stigma, inflexible scheduling, and uncertain eligibility. For immigrants, these obstacles may be compounded by migration status, prior experiences of state authority, and fear that personal information could be shared beyond the clinical setting. A systematic review of immigrant mental-health service use in the United States found lower utilization among several immigrant groups, with language barriers, stigma, limited service knowledge, and culturally incongruent care contributing to unmet need (Derr, 2016).

Trust is particularly consequential in trauma-informed practice because engagement depends upon confidence that disclosure will not produce humiliation, coercion, discrimination, or institutional harm. Refugee women in high-income resettlement countries report barriers involving stigmatization, language, limited culturally appropriate resources, socioeconomic constraints, and concerns surrounding service acceptability, while awareness and social support can facilitate access (DeSa et al., 2022). For trauma-exposed clients, mistrust should not

automatically be interpreted as resistance; it may reflect prior persecution, exclusion, gendered violence, discriminatory treatment, or unfamiliar institutions.

Institutional mistrust becomes acute where healthcare systems are experienced as extensions of immigration control or other coercive authorities. Research in United Kingdom immigration removal centres demonstrates that stigma, powerlessness, confusion over entitlement, and conflictual intergroup relationships can undermine healthcare trust and contribute to disengagement, even when services are formally available (Kéllezi et al., 2021). Urban behavioral health organizations serving immigrant populations must consequently separate clinical care visibly from enforcement functions, communicate confidentiality boundaries clearly, and ensure administrative procedures do not reproduce surveillance or threat.

African evidence illustrates how exclusion can be embedded in ordinary service encounters. Among Zimbabwean migrants seeking public healthcare in South Africa, discriminatory practices based on nationality were documented within institutions responsible for providing care, demonstrating how xenophobia can transform formal entitlement into unequal access (Crush & Tawodzera, 2014). Reports of disrespect or exclusion can circulate through migrant networks, shaping collective expectations about whether institutions are trustworthy.

Structural barriers also affect non-migrant populations and may shape expectations of care after migration. In Nigeria's Niger Delta, service users identified financial cost, geographic distance, limited knowledge, waiting times, and stigma as barriers to sustained mental-health service utilization (Jack-Ide & Uys, 2013). Immigrants may therefore arrive with experiences of inaccessible systems that influence expectations of behavioral healthcare in destination cities.

Access is also shaped by developmental and generational context. A systematic scoping review of immigrant students in high-income countries identified stigma, limited mental-health knowledge, cultural expectations, and service-navigation difficulties among important barriers, while mental-health literacy and adaptation could facilitate utilization (Dombou et al., 2023). First-generation young people may simultaneously negotiate parental attitudes toward treatment, confidentiality concerns, financial dependence, and institutional unfamiliarity.

Culturally responsive trauma-informed systems should therefore conceptualize trust as an organizational outcome rather than a personality trait of the client. Accessible care requires multilingual navigation, eligibility rules, pathways, continuity with clinicians, community partnerships, and safeguards against discriminatory treatment. Staff should explain confidentiality, consent, documentation requirements, referral processes, and information-sharing limits in plain language. Organizations should examine whether intake practices, security procedures, billing arrangements, interpreter access, or identification requirements create avoidable deterrents. Community advisory structures and service-user feedback can identify barriers invisible to administrators, while partnerships with immigrant-led organizations can strengthen credibility. Access is achieved when services are not merely present, but practically usable, culturally intelligible, relationally safe, and institutionally worthy of trust.

4.2 Workforce Competence and Organizational Readiness

Workforce competence and organizational readiness are decisive conditions for translating culturally responsive, trauma-informed principles into routine behavioral health practice. In urban services serving immigrant and first-generation populations, competence requires more than technical knowledge of trauma. Staff must recognize culturally patterned expressions of distress, communicate across linguistic differences, understand migration-related stressors,

manage implicit bias, and respond without reproducing coercion or stereotyping. South African evidence indicates that health professionals identify cultural and linguistic competence as areas requiring stronger curricular preparation and continuing professional development in diverse service environments (Matthews & Van Wyk, 2018).

Organizational readiness extends individual capabilities into governance, workflows, supervision, and resource allocation. Trauma-informed implementation is more sustainable when training is embedded within broader organizational change rather than delivered as an isolated educational event. A systematic review found that staff-focused trauma-informed interventions commonly improved knowledge, attitudes, and some practice behaviors, although evidence quality varied (Purtle, 2020). Implementation research likewise identifies leadership support, resources, staff attitudes, interdisciplinary collaboration, and organizational culture as recurring enablers or barriers to trauma-informed care (Huo et al., 2023).

Multidisciplinary competence is especially important when clients present with interacting psychological, medical, social, linguistic, and legal needs. Evidence from emergency-care settings demonstrates that coordinated multidisciplinary practice can strengthen communication, decision-making, efficiency, and patient-centered outcomes, principles relevant to complex urban behavioral health systems (Ohunyon, Ayinde & Ojukwu, 2023). Workforce effectiveness also depends on professional attributes such as communication, teamwork, reliability, and interpersonal competence, which have been associated with organizational performance in Nigerian institutions (Onche, Adenekan & Sabboh, 2021).

Readiness additionally requires mechanisms for developing and retaining capable personnel. Organizational models of talent development emphasize continuous learning, competency mapping, and alignment between workforce capabilities and strategic objectives, offering useful principles for complex service environments (Falemi, Akhigbe & Akin-Oluyomi, 2020). Retention is equally consequential because culturally responsive trauma care depends upon continuity, accumulated expertise, and stable therapeutic relationships; workforce frameworks linking employee retention with operational performance reinforce the value of treating staffing stability strategically (Falemi, Akhigbe & Akin-Oluyomi, 2023).

5. Practice, Policy, and Research Priorities

5.1 Advancing Equitable Trauma-Informed Behavioral Health

Advancing equitable trauma-informed behavioral health requires a shift from episodic, diagnosis-centred provision toward systems that recognize how trauma, migration, culture, race, language, socioeconomic position, and legal status interact to shape access and recovery. Equity does not mean providing identical services to every client; it requires proportionate responses to unequal barriers and differing needs. For immigrant and first-generation populations in urban settings, this entails designing services that are clinically competent, culturally intelligible, accessible, and flexible enough to address structural stressors alongside psychological symptoms. Reviews of refugee mental health emphasize that culturally responsive and trauma-informed care must account for barriers such as discrimination, legal insecurity, language discordance, stigma, financial hardship, and limited availability of appropriately trained professionals (Mattar & Gellatly, 2022).

A central priority is to move trauma-informed care beyond individual clinician sensitivity and embed it within organizational structures. Refugee-serving professionals have reported inconsistencies in translating trauma-informed principles into resettlement practice, particularly

where cultural adaptation, interpreter collaboration, cross-sector communication, and community participation are weak. Building equitable systems therefore requires protocols, sustained workforce development, interagency learning, and partnerships with communities rather than single cultural-competence training (Im & Swan, 2021). Such organizational reform is especially important in urban behavioral health systems, where fragmented pathways can force clients to repeat traumatic experiences while navigating disconnected agencies.

Equity also requires a continuum of support rather than reliance on specialist psychotherapy alone. Refugee mental-health models increasingly advocate layered services that connect basic safety and social supports with prevention, family and community interventions, and specialized clinical treatment according to need. This approach recognizes that distress may be sustained by housing instability, employment barriers, family separation, discrimination, or resettlement difficulties that cannot be resolved through symptom-focused care alone (Bunn et al., 2023). For first-generation clients, prevention should address intergenerational strain, identity negotiation, school or workplace discrimination, and family communication before problems escalate into crisis. Policy environments can either protect or undermine trauma-informed practice. Australian evidence concerning refugees and asylum seekers with insecure status illustrates how legal uncertainty, economic exclusion, and restricted access to supports can intensify psychological distress, making policy itself a determinant of mental-health risk (Kenny, Grech & Procter, 2022). Equitable behavioral health therefore requires coordination between clinical systems and social-policy structures, while ensuring that health services remain trusted spaces rather than perceived extensions of immigration enforcement or other coercive institutions.

Workforce capacity must also be strengthened. European evidence shows that professionals caring for migrants and refugees encounter language barriers, unfamiliar explanatory models, complex social needs, limited resources, and uncertainty about culturally appropriate practice, all of which can constrain therapeutic engagement (Peñuela-O'Brien et al., 2023). Urban systems should consequently invest in qualified interpreters, reflective supervision, cultural humility, trauma competencies, anti-discrimination practice, and recruitment of clinicians and community workers who reflect the populations served.

Low- and middle-income settings further demonstrate why equity requires health-system strengthening rather than isolated clinical innovation. In Nigeria, challenges in financing, legislation, workforce development, research, and integration of mental health into primary care have constrained accessible and affordable services, reinforcing the importance of sustained policy implementation and public investment (Wada et al., 2021). Ghanaian evidence similarly indicates that health-system structure and service processes influence consumers' perceptions of mental-health quality, suggesting that availability, accessibility, professional competence, therapeutic relationships, and user experience should be treated as interconnected dimensions of equity (Badu et al., 2020). Equitable improvement also depends on accountable use of operational information. Conceptual work linking workforce behavior with real-time service data demonstrates how organizations can identify bottlenecks, monitor responsiveness, and adapt resource allocation more rapidly (Akhigbe, Falemi & Akin-Oluyomi, 2023). In behavioral health, analogous systems should be governed ethically and used to monitor disparities rather than intensify surveillance. Organizations can stratify access, waiting times, dropout, interpreter use, crisis utilization, and experience by language, migration generation, and other relevant characteristics while protecting confidentiality. These data should be combined with community advisory input because numerical indicators alone cannot reveal whether care feels respectful,

safe, or culturally meaningful. Advancing equity ultimately requires institutional accountability: leadership must link trauma-informed commitments to measurable access, experience, continuity, workforce, and outcome indicators, examine disparities, and redesign services with immigrant and first-generation communities as partners in determining what recovery-oriented care should look like.

Conclusion

This review has demonstrated that culturally responsive trauma-informed behavioral health requires more than the application of universal clinical principles. Its central aim was to examine how migration, first-generation identity, cultural meaning, family relationships, language, structural inequality, and organizational conditions shape the experience of trauma and the effectiveness of care in urban settings. Across the reviewed evidence, this objective was met by showing that trauma is interpreted through cultural, social, and intergenerational frameworks, while recovery is influenced by trust, belonging, communication, family support, community resources, and access to responsive institutions.

The review further established that effective assessment depends on cultural humility, valid and context-sensitive measures, careful interpretation of idioms of distress, and attention to ongoing post-migration stressors. Language access emerged as a core clinical requirement rather than an administrative convenience, particularly where interpreter quality, confidentiality, and conceptual equivalence affect disclosure and therapeutic alliance. Family and community participation were also shown to be important when guided by client preference, safety, and respect for cultural and spiritual practices. At the systems level, workforce competence, multidisciplinary collaboration, organizational readiness, and institutional trust were identified as essential determinants of equitable service delivery.

These findings support the conclusion that trauma-informed care becomes responsive only when cultural and structural realities are incorporated into assessment, engagement, treatment, and service design. Urban behavioral health systems should therefore invest in qualified interpreters, culturally informed workforce development, reflective supervision, community partnerships, and mechanisms for monitoring disparities in access, continuity, and outcomes. Policy should strengthen confidentiality protections, reduce administrative and financial barriers, and support services that remain clearly separated from coercive or enforcement functions. Future research should prioritize longitudinal, community-engaged, and culturally validated studies involving diverse immigrant and first-generation populations. Greater attention should also be given to intersectionality, intergenerational processes, indigenous and community-based healing, and implementation strategies that translate cultural responsiveness from principle into sustainable organizational practice.

References

- Abdi, S.M., Miller, A.B., Agalab, N.Y. and Ellis, B.H. (2022) 'Partnering with refugee communities to improve mental health access: Going from "why are they not coming" to "what can I (we) do differently?"', *Cultural Diversity and Ethnic Minority Psychology*, 28(3), pp. 370–378. DOI: [10.1037/cdp0000476](https://doi.org/10.1037/cdp0000476).
- Aggarwal, N.K., Jarvis, G.E., Gómez-Carrillo, A., Kirmayer, L.J. and Lewis-Fernández, R. (2020) 'The Cultural Formulation Interview since DSM-5: Prospects for training, research, and clinical practice', *Transcultural Psychiatry*, 57(4), pp. 496–514. DOI: [10.1177/1363461520940481](https://doi.org/10.1177/1363461520940481).
- Akhigbe, R., Falemi, A. and Akin-Oluyomi, O.T. (2023) 'A conceptual model for improving customer experience using workforce behavior and real-time operational data', *International Journal of Multidisciplinary Research and Growth Evaluation*, 4(6), pp. 1322–1338. DOI: [10.54660/IJMRGE.2023.4.6.1322-1338](https://doi.org/10.54660/IJMRGE.2023.4.6.1322-1338).
- Badu, E., O'Brien, A.P., Mitchell, R. and Osei, A. (2020) 'Mediation and moderation effects of health system structure and process on the quality of mental health services in Ghana – structural equation modelling', *PLOS ONE*, 15(5), e0233351. DOI: [10.1371/journal.pone.0233351](https://doi.org/10.1371/journal.pone.0233351).
- Balogun, C.T. and Sabboh, G.M. (2021) 'Psycho-sociological factors as determinants of juvenile delinquency among adolescents in junior secondary schools in Patani Local Government Area of Delta State, Nigeria', *International Journal of Academic Management Science Research*, 5(10), pp. 34–40.
- Bauer, A.M. and Alegria, M. (2010) 'Impact of patient language proficiency and interpreter service use on the quality of psychiatric care: A systematic review', *Psychiatric Services*, 61(8), pp. 765–773. DOI: [10.1176/ps.2010.61.8.765](https://doi.org/10.1176/ps.2010.61.8.765).
- Blackmore, R., Boyle, J.A., Fazel, M., Ranasinha, S., Gray, K.M., Fitzgerald, G., Misso, M. and Gibson-Helm, M. (2020) 'The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis', *PLOS Medicine*, 17(9), e1003337. DOI: [10.1371/journal.pmed.1003337](https://doi.org/10.1371/journal.pmed.1003337).
- Bunn, M., Khanna, D., Farmer, E., Esbrook, E., Ellis, H., Richard, A. and Weine, S. (2023) 'Rethinking mental healthcare for refugees', *SSM – Mental Health*, 3, 100196. DOI: [10.1016/j.ssmmh.2023.100196](https://doi.org/10.1016/j.ssmmh.2023.100196).
- Cleary, M. and Hungerford, C. (2015) 'Trauma-informed care and the research literature: How can the mental health nurse take the lead to support women who have survived sexual assault?', *Issues in Mental Health Nursing*, 36(5), pp. 370–378. DOI: [10.3109/01612840.2015.1009661](https://doi.org/10.3109/01612840.2015.1009661).
- Crush, J. and Tawodzera, G. (2014) 'Medical xenophobia and Zimbabwean migrant access to public health services in South Africa', *Journal of Ethnic and Migration Studies*, 40(4), pp. 655–670. DOI: [10.1080/1369183X.2013.830504](https://doi.org/10.1080/1369183X.2013.830504).
- Derr, A.S. (2016) 'Mental health service use among immigrants in the United States: A systematic review', *Psychiatric Services*, 67(3), pp. 265–274. DOI: [10.1176/appi.ps.201500004](https://doi.org/10.1176/appi.ps.201500004).
- DeSa, S., Gebremeskel, A.T., Omonaiye, O. and Yaya, S. (2022) 'Barriers and facilitators to access mental health services among refugee women in high-income countries: A systematic review', *Systematic Reviews*, 11, Article 62. DOI: [10.1186/s13643-022-01936-1](https://doi.org/10.1186/s13643-022-01936-1).

- Diri, B.H. (2023) 'Language barriers: An overview for efficiency in Nigeria's health system', *Journal of Health, Applied Sciences and Management*, 7(1), pp. 85–90. DOI: [10.4314/johasam.v7i1.9](https://doi.org/10.4314/johasam.v7i1.9).
- Dombou, C., Omonaiye, O., Fraser, S., Cénat, J.M., Fournier, K. and Yaya, S. (2023) 'Barriers and facilitators associated with the use of mental health services among immigrant students in high-income countries: A systematic scoping review', *PLOS ONE*, 18(6), e0287162. DOI: [10.1371/journal.pone.0287162](https://doi.org/10.1371/journal.pone.0287162).
- Escamilla, S. and Saasa, S. (2020) 'Gendered patterns in depression and anxiety among African immigrants in the United States', *Journal of Evidence-Based Social Work*, 17(4), pp. 392–405. DOI: [10.1080/26408066.2020.1748153](https://doi.org/10.1080/26408066.2020.1748153).
- Ezeani, P.T. and Sabboh, G.M. (2021) 'Psychosocial predictors of the changing role of husbands as breadwinners in a depressed economy in Ibadan Metropolis, Oyo State, Nigeria', *British Journal of Education, Learning and Development Psychology*, 4(2), pp. 31–42. DOI: [10.52589/BJELDP-ZU3KSUJU](https://doi.org/10.52589/BJELDP-ZU3KSUJU).
- Ezeani, P.T., Ogu, C. and Sabboh, G.M. (2023) 'Parental involvement as correlate of career decision-making among secondary school students in Oyo State, Nigeria', *British Journal of Multidisciplinary and Advanced Studies: Arts, Humanities and Social Sciences*, 4(4), pp. 106–115. DOI: [10.37745/bjmas.2022.0260](https://doi.org/10.37745/bjmas.2022.0260).
- Falemi, A., Akhigbe, R. and Akin-Oluyomi, O.T. (2020) 'A conceptual supply chain talent development model for capability building across distributed operations', *International Journal of Multidisciplinary Research and Growth Evaluation*, 1(5), pp. 439–456. DOI: [10.54660/IJMRGE.2020.1.5.439-456](https://doi.org/10.54660/IJMRGE.2020.1.5.439-456).
- Falemi, A., Akhigbe, R. and Akin-Oluyomi, O.T. (2023) 'A proposed framework for integrating employee retention systems with operational productivity metrics', *International Journal of Advanced Multidisciplinary Research and Studies*, 3(1), pp. 1704–1724. DOI: [10.62225/2583049X.2023.3.1.5339](https://doi.org/10.62225/2583049X.2023.3.1.5339).
- Fennig, M. and Denov, M. (2021) 'Interpreters working in mental health settings with refugees: An interdisciplinary scoping review', *American Journal of Orthopsychiatry*, 91(1), pp. 50–65. DOI: [10.1037/ort0000518](https://doi.org/10.1037/ort0000518).
- Fortuna, L.R., Porche, M.V. and Alegría, M. (2008) 'Political violence, psychosocial trauma, and the context of mental health services use among immigrant Latinos in the United States', *Ethnicity & Health*, 13(5), pp. 435–463. DOI: [10.1080/13557850701837286](https://doi.org/10.1080/13557850701837286).
- Hagan, S., Hunt, X., Kilian, S., Chiliza, B. and Swartz, L. (2020) 'Ad hoc interpreters in South African psychiatric services: Service provider perspectives', *Global Health Action*, 13(1), Article 1684072. DOI: [10.1080/16549716.2019.1684072](https://doi.org/10.1080/16549716.2019.1684072).
- Hook, J.N., Davis, D.E., Owen, J., Worthington, E.L. Jr. and Utsey, S.O. (2013) 'Cultural humility: Measuring openness to culturally diverse clients', *Journal of Counseling Psychology*, 60(3), pp. 353–366. DOI: [10.1037/a0032595](https://doi.org/10.1037/a0032595).
- Huo, Y., Couzner, L., Windsor, T., Laver, K., Dissanayaka, N.N. and Cations, M. (2023) 'Barriers and enablers for the implementation of trauma-informed care in healthcare settings: A systematic review', *Implementation Science Communications*, 4, Article 49. DOI: [10.1186/s43058-023-00428-0](https://doi.org/10.1186/s43058-023-00428-0).
- Hynie, M. (2018) 'The social determinants of refugee mental health in the post-migration context: A critical review', *The Canadian Journal of Psychiatry*, 63(5), pp. 297–303. DOI: [10.1177/0706743717746666](https://doi.org/10.1177/0706743717746666).

- Im, H. and Swan, L.E.T. (2021) 'Working towards culturally responsive trauma-informed care in the refugee resettlement process: Qualitative inquiry with refugee-serving professionals in the United States', *Behavioral Sciences*, 11(11), Article 155. DOI: [10.3390/bs11110155](https://doi.org/10.3390/bs11110155).
- Im, H., Ferguson, A. and Hunter, M. (2017) 'Cultural translation of refugee trauma: Cultural idioms of distress among Somali refugees in displacement', *Transcultural Psychiatry*, 54(5–6), pp. 626–652. DOI: [10.1177/1363461517744989](https://doi.org/10.1177/1363461517744989).
- Im, H., Rodriguez, C. and Grumbine, J.M. (2021) 'A multitier model of refugee mental health and psychosocial support in resettlement: Toward trauma-informed and culture-informed systems of care', *Psychological Services*, 18(3), pp. 345–364. DOI: [10.1037/ser0000412](https://doi.org/10.1037/ser0000412).
- Jack-Ide, I.O. and Uys, L. (2013) 'Barriers to mental health services utilization in the Niger Delta region of Nigeria: Service users' perspectives', *Pan African Medical Journal*, 14, Article 159. DOI: [10.11604/pamj.2013.14.159.1970](https://doi.org/10.11604/pamj.2013.14.159.1970).
- Këllezi, B., Wakefield, J., Bowe, M., Stevenson, C. and McNamara, N. (2021) 'Healthcare provision inside immigration removal centres: A social identity analysis of trust, legitimacy and disengagement', *Applied Psychology: Health and Well-Being*, 13(3), pp. 578–601. DOI: [10.1111/aphw.12263](https://doi.org/10.1111/aphw.12263).
- Kenny, M.A., Grech, C. and Procter, N. (2022) 'A trauma-informed response to COVID-19 and the deteriorating mental health of refugees and asylum seekers with insecure status in Australia', *International Journal of Mental Health Nursing*, 31(1), pp. 62–69. DOI: [10.1111/inm.12932](https://doi.org/10.1111/inm.12932).
- Kern, M.R., Duinhof, E.L., Walsh, S.D., Cosma, A., Moreno-Maldonado, C., Molcho, M., Currie, C. and Stevens, G.W.J.M. (2020) 'Intersectionality and adolescent mental well-being: A cross-nationally comparative analysis of the interplay between immigration background, socioeconomic status and gender', *Journal of Adolescent Health*, 66(6S), pp. S12–S20. DOI: [10.1016/j.jadohealth.2020.02.013](https://doi.org/10.1016/j.jadohealth.2020.02.013).
- Kirmayer, L.J., Narasiah, L., Munoz, M., Rashid, M., Ryder, A.G., Guzder, J., Hassan, G., Rousseau, C. and Pottie, K. (2011) 'Common mental health problems in immigrants and refugees: General approach in primary care', *CMAJ*, 183(12), pp. E959–E967. DOI: [10.1503/cmaj.090292](https://doi.org/10.1503/cmaj.090292).
- Kpobi, L.N.A. and Swartz, L. (2019) 'Muslim traditional healers in Accra, Ghana: Beliefs about and treatment of mental disorders', *Journal of Religion and Health*, 58(3), pp. 833–846. DOI: [10.1007/s10943-018-0668-1](https://doi.org/10.1007/s10943-018-0668-1).
- Lewis-Fernández, R. and Kirmayer, L.J. (2019) 'Cultural concepts of distress and psychiatric disorders: Understanding symptom experience and expression in context', *Transcultural Psychiatry*, 56(4), pp. 786–803. DOI: [10.1177/1363461519861795](https://doi.org/10.1177/1363461519861795).
- Li, S.S.Y., Liddell, B.J. and Nickerson, A. (2016) 'The relationship between post-migration stress and psychological disorders in refugees and asylum seekers', *Current Psychiatry Reports*, 18(9), Article 82. DOI: [10.1007/s11920-016-0723-0](https://doi.org/10.1007/s11920-016-0723-0).
- Madigoe, T., Burns, J., Zhang, M. and Subramaney, U. (2017) 'Towards a culturally appropriate trauma assessment in a South African Zulu community', *Psychological Trauma: Theory, Research, Practice, and Policy*, 9(3), pp. 274–281. DOI: [10.1037/tra0000231](https://doi.org/10.1037/tra0000231).
- Mak, C. and Wieling, E. (2022) 'A systematic review of evidence-based family interventions for trauma-affected refugees', *International Journal of Environmental Research and Public Health*, 19(15), Article 9361. DOI: [10.3390/ijerph19159361](https://doi.org/10.3390/ijerph19159361).

- Mancenido, A., Williams, E.C. and Hajat, A. (2020) 'Examining psychological distress across intersections of immigrant generational status, race, poverty, and gender', *Community Mental Health Journal*, 56(7), pp. 1269–1274. DOI: [10.1007/s10597-020-00584-w](https://doi.org/10.1007/s10597-020-00584-w).
- Mattar, S. and Gellatly, R. (2022) 'Refugee mental health: Culturally relevant considerations', *Current Opinion in Psychology*, 47, 101429. DOI: [10.1016/j.copsyc.2022.101429](https://doi.org/10.1016/j.copsyc.2022.101429).
- Matthews, M. and Van Wyk, J. (2018) 'Towards a culturally competent health professional: A South African case study', *BMC Medical Education*, 18, Article 112. DOI: [10.1186/s12909-018-1187-1](https://doi.org/10.1186/s12909-018-1187-1).
- Mbazzi, F.B., Dewailly, A., Admasu, K., Duagani, Y., Wamala, K., Vera, A., Bwesigye, D. and Roth, G. (2021) 'Cultural adaptations of the standard EMDR protocol in five African countries', *Journal of EMDR Practice and Research*, 15(1), pp. 29–43. DOI: [10.1891/EMDR-D-20-00028](https://doi.org/10.1891/EMDR-D-20-00028).
- Ohunyon, G., Ayinde, M. and Ojukwu, P.U. (2023) 'The impact of multidisciplinary collaboration on patient outcomes in emergency departments', *International Journal of Advanced Multidisciplinary Research and Studies*, 3(6), pp. 2727–2741. DOI: [10.62225/2583049X.2023.3.6.6029](https://doi.org/10.62225/2583049X.2023.3.6.6029).
- Onche, V.O., Adenekan, E.T. and Sabboh, G.M. (2021) 'Influence of personal and professional attributes of secretaries on organisational success of universities in Oyo State, Nigeria', *International Journal of Science Academic Research*, 2(4), pp. 1355–1359.
- Oyarekua, T.A. and Sabboh, G.M. (2021) 'Parenting styles and effortful control as predictors of mental health among secondary school adolescents in Akinyele Local Government Area, Oyo State, Nigeria', *International Journal of Academic Multidisciplinary Research*, 5(10), pp. 108–114.
- Patel, A.R. and Hall, B.J. (2021) 'Beyond the DSM-5 diagnoses: A cross-cultural approach to assessing trauma reactions', *Focus*, 19(2), pp. 197–203. DOI: [10.1176/appi.focus.20200049](https://doi.org/10.1176/appi.focus.20200049).
- Peñuela-O'Brien, E., Wan, M.W., Edge, D. and Berry, K. (2023) 'Health professionals' experiences of and attitudes towards mental healthcare for migrants and refugees in Europe: A qualitative systematic review', *Transcultural Psychiatry*, 60(1), pp. 176–198. DOI: [10.1177/13634615211067360](https://doi.org/10.1177/13634615211067360).
- Pittaway, T., Riggs, E. and Dantas, J.A.R. (2023) 'Intergenerational conflict among resettled South Sudanese in Australia', *Transcultural Psychiatry*, 60(2), pp. 215–225. DOI: [10.1177/13634615221105115](https://doi.org/10.1177/13634615221105115).
- Purtle, J. (2020) 'Systematic review of evaluations of trauma-informed organizational interventions that include staff trainings', *Trauma, Violence, & Abuse*, 21(4), pp. 725–740. DOI: [10.1177/1524838018791304](https://doi.org/10.1177/1524838018791304).
- Raghavan, S.S. (2019) 'Cultural considerations in the assessment of survivors of torture', *Journal of Immigrant and Minority Health*, 21(3), pp. 586–595. DOI: [10.1007/s10903-018-0787-5](https://doi.org/10.1007/s10903-018-0787-5).
- Reeves, E. (2015) 'A synthesis of the literature on trauma-informed care', *Issues in Mental Health Nursing*, 36(9), pp. 698–709. DOI: [10.3109/01612840.2015.1025319](https://doi.org/10.3109/01612840.2015.1025319).
- Renzaho, A.M.N., Dhingra, N. and Georgeou, N. (2017) 'Youth as contested sites of culture: The intergenerational acculturation gap amongst new migrant communities—Parental and young adult perspectives', *PLOS ONE*, 12(2), e0170700. DOI: [10.1371/journal.pone.0170700](https://doi.org/10.1371/journal.pone.0170700).

- Sabboh, G.M. (2021) 'Psychosocial factors as predictors of moral disengagement among in-school adolescents in Ibadan Metropolis', *International Journal of Academic Information Systems Research*, 5(10), pp. 1–7.
- Sabboh, G.M. (2022) Development and validation of the revised self-esteem scale for Nigerian adolescents and youths. Master's thesis. Moscow, Russia: National Research University Higher School of Economics. DOI: [10.13140/RG.2.2.32753.45926](https://doi.org/10.13140/RG.2.2.32753.45926).
- Sabboh, G.M. (2023) Relationship of perceived stress, hopelessness and self-esteem with attitude towards suicide among people in Nigeria. Master's thesis. Moscow, Russia: National Research University Higher School of Economics. DOI: [10.13140/RG.2.2.21009.40802](https://doi.org/10.13140/RG.2.2.21009.40802).
- Sabboh, G.M. and Alausa, S.S. (2021) 'School climate, self-efficacy, social media and peer pressure as correlates of aggressive behaviour among secondary school adolescents in Ibadan Metropolis, Oyo State, Nigeria', *International Journal of Academic Management Science Research*, 5(11), pp. 43–51.
- Sabboh, G.M. and Ojo, Y. (2023) 'Influence of peer approval seeking, socio-economic status and locus of control on academic self-handicapping behaviour of undergraduates in Oyo State, Nigeria', *British Journal of Multidisciplinary and Advanced Studies: Education, Learning, Training & Development*, 4(4), pp. 98–112. DOI: [10.37745/bjmas.2022.0264](https://doi.org/10.37745/bjmas.2022.0264).
- Sabboh, G.M. and Olugbade, D. (2021) 'Influence of teaching styles and parental bonding on academic boredom among undergraduates in Oyo State, Nigeria', *African Journal of Social Sciences and Humanities Research*, 4(1), pp. 122–133.
- Sabboh, G.M. and Yusuff, A.O. (2021) 'Influence of psycho-social factors as predictors of child abuse among junior secondary school students in Kaduna Metropolis of Kaduna State', *International Journal of Academic and Applied Research*, 5(8), pp. 18–24.
- Schlechter, P., Wilkinson, P.O., Knausenberger, J., Wanninger, K., Kamp, S., Morina, N. and Hellmann, J.H. (2021) 'Depressive and anxiety symptoms in refugees: Insights from classical test theory, item response theory and network analysis', *Clinical Psychology & Psychotherapy*, 28(1), pp. 169–181. DOI: [10.1002/cpp.2499](https://doi.org/10.1002/cpp.2499).
- Sweeney, A., Clement, S., Filson, B. and Kennedy, A. (2016) 'Trauma-informed mental healthcare in the UK: What is it and how can we further its development?', *Mental Health Review Journal*, 21(3), pp. 174–192. DOI: [10.1108/MHRJ-01-2015-0006](https://doi.org/10.1108/MHRJ-01-2015-0006).
- Thela, L., Tomita, A., Maharaj, V., Mhlongo, M. and Burns, J.K. (2017) 'Counting the cost of Afrophobia: Post-migration adaptation and mental health challenges of African refugees and migrants in South Africa', *Transcultural Psychiatry*, 54(5–6), pp. 715–732. DOI: [10.1177/1363461517745472](https://doi.org/10.1177/1363461517745472).
- Tujague, N.A. and Ryan, K.L. (2021) 'Ticking the box of “cultural safety” is not enough: Why trauma-informed practice is critical to Indigenous healing', *Rural and Remote Health*, 21(3), Article 6411. DOI: [10.22605/RRH6411](https://doi.org/10.22605/RRH6411).
- Viruell-Fuentes, E.A., Miranda, P.Y. and Abdulrahim, S. (2012) 'More than culture: Structural racism, intersectionality theory, and immigrant health', *Social Science & Medicine*, 75(12), pp. 2099–2106. DOI: [10.1016/j.socscimed.2011.12.037](https://doi.org/10.1016/j.socscimed.2011.12.037).
- Wada, Y.H., Rajwani, L., Anyam, E., Karikari, E., Njikizana, M., Srour, L. and Khalid, G.M. (2021) 'Mental health in Nigeria: A neglected issue in public health', *Public Health in Practice*, 2, 100166. DOI: [10.1016/j.puhip.2021.100166](https://doi.org/10.1016/j.puhip.2021.100166).
- Walker, R. and Vearey, J. (2023) ““Let’s manage the stressor today”: Exploring the mental health response to forced migrants in Johannesburg, South Africa”, *International Journal*

- of Migration, Health and Social Care, 19(1), pp. 1–15. DOI: [10.1108/IJMHSC-11-2021-0103](https://doi.org/10.1108/IJMHSC-11-2021-0103).
- Wyatt, G.E., Thames, A., Simbayi, L., Stein, D.J., Burns, J. and Maselesele, M. (2017) ‘Trauma and mental health in South Africa: Overview’, *Psychological Trauma: Theory, Research, Practice, and Policy*, 9(3), pp. 249–251. DOI: [10.1037/tra0000144](https://doi.org/10.1037/tra0000144).
- Wylie, L., Van Meyel, R., Harder, H., Sukhera, J., Luc, C., Ganjavi, H., Elfakhani, M. and Wardrop, N. (2018) ‘Assessing trauma in a transcultural context: Challenges in mental health care with immigrants and refugees’, *Public Health Reviews*, 39, Article 22. DOI: [10.1186/s40985-018-0102-y](https://doi.org/10.1186/s40985-018-0102-y).
- Yusuf, A.O. and Sabboh, G.M. (2021) ‘Psychological factors as correlates of suicidal ideation among university students in Lagos State’, *International Journal of Academic and Applied Research*, 5(9), pp. 1–7.
- Yusuf, A.O., Sabboh, G.M. and Asiyambi, M. (2023) ‘Correlates of academic buoyancy among undergraduates in the Ibadan Metropolis’, *International Journal of Academic Information Systems Research*, 7(1), pp. 1–10.